

The Euler-characteristic of discrete groups and Yuzvinskii's entropy addition formula

Gábor Elek

February 1, 2008

ABSTRACT: We prove that Yuzvinskii's entropy addition formula can not be extended for the action of certain finitely generated nonamenable groups.

1991 AMS CLASSIFICATION NUMBERS: 22D40, 58F03

KEYWORDS: entropy, groups of finite type, symbolic dynamics

Gábor Elek

Mathematical Institute, Hungarian Academy of Sciences

Budapest 1364, POB 127 Hungary

e-mail elek@math-inst.hu

1 Introduction

Yuzvinskii's entropy addition formula [7],[9]

If α is a $\mathbf{Z}^d$ -action on the compact group X , and Y is an α -invariant normal subgroup of X , let α_Y denote the restriction of α to Y and $\alpha_{X|Y}$ be the induced action on X/Y . Then

$$h(\alpha) = h(\alpha_{X|Y}) + h(\alpha_Y) \quad ,$$

where $h(\alpha)$ is the topological $\mathbf{Z}^d$ -entropy of α .

Historically, the notion of topological entropy was introduced for $\mathbf{Z}$ -actions and $\mathbf{Z}^d$ -actions. Later Kieffer [6] extended the definition for amenable groups. Recently, Friedland introduced a very general definition of entropy for arbitrary finitely generated group and semigroup actions [5]. Although the addition formula is not extended yet for finitely generated amenable groups in general, the work of Ward and Zhang [8] suggests that such extension is possible. We shall show in this paper that although Friedland's notion of entropy is very nice and useful, it cannot be used to extend the addition formula for arbitrary groups and semigroups. In fact, there is no sensible notion of entropy for which the addition formula can be extended for free groups, even if we consider group actions only on compact, metrizable, Abelian groups.

2 The axioms for the addition formula in symbolic dynamics

Let Γ be a countable group or semigroup and let S be a finite abelian group. Then we consider as usual the $\Sigma(\Gamma, S)$ Bernoulli-shift [4] of S -valued functions on Γ . Then $\Sigma(\Gamma, S)$ is a compact, metrizable, Abelian group with a natural Γ -action, where the action on the left is given by

$$\sigma_\gamma(\tau)(\gamma') = \tau(\gamma'\gamma) \quad .$$

A linear subshift is a Γ -invariant closed subgroup of $\Sigma(\Gamma, S)$. Below we give a system of axioms for the notion of entropy. We call a group or a semigroup *SEA* (symbolically

entropy additive) if the following axioms hold:

- (A1) There is a real valued entropy function h on linear subshifts.
- (A2) $h(\sum(\Gamma, S)) = \log_2 |S|$ for any finite Abelian group S .
- (A3) If $\phi_1 \subset \phi_2$ are linear subshifts then $0 \leq h(\phi_1) \leq h(\phi_2)$.
- (A4) If $\phi_1 \subset \sum(\Gamma, S_1), \phi_2 \subset \sum(\Gamma, S_2)$ and $T : \phi_1 \rightarrow \phi_2$ is a Γ -invariant homeomorphism, then $h(\phi_1) = h(\phi_2)$.
- (A5) If a linear subshift ϕ contains only finite many elements, then $h(\phi) = 0$.
- (A6) (The Yuzvinskii-axiom) If $T : \phi_1 \rightarrow \phi_2$ is a Γ -invariant surjective continuous map between linear subshifts, then

$$h(Ker \phi_1) + h(\phi_2) = h(\phi_1) \quad .$$

Note that the notions of entropy mentioned in the Introduction satisfy the first five axioms. So far we know that $\mathbf{N}, \mathbf{Z}$ and $\mathbf{Z}^d$ are *SEA*. Before stating our main result we prove two easy propositions.

Proposition 2.1 *If Γ_1 is a quotient group of Γ and Γ_1 is not SEA, then Γ is not SEA either.*

Proof: Let $\pi : \Gamma \rightarrow \Gamma_1$ be a surjective group homomorphism. Then for any finite Abelian group S , we have a natural injective map

$$\pi^* : \sum(\Gamma_1, S) \rightarrow \sum(\Gamma, S)$$

defined by $\pi^*(\tau)(\gamma) = \tau(\pi(\gamma))$. Now let us notice that for any $\gamma \in \Gamma$,

$$\pi^*(\sigma_{\pi(\gamma)}(\tau)) = \sigma_\gamma(\pi^*(\tau)) \quad .$$

Indeed,

$$\pi^*(\sigma_{\pi(\gamma)}(\tau))(\gamma') = \sigma_{\pi(\gamma)}(\tau)(\pi(\gamma')) = \tau(\pi(\gamma'\gamma)) = \pi^*(\tau)(\gamma'\gamma) = \sigma_\gamma(\pi^*(\tau))(\gamma') \quad .$$

Consequently, by continuity, π^* maps linear subshifts into linear subshifts. Let $T : \phi_1 \rightarrow \phi_2$ be a continuous group homomorphism between linear Γ_1 -subshifts. Then we can define

$$\pi^*T : \pi^*\phi_1 \rightarrow \pi^*\phi_2 \quad ,$$

by

$$(\pi^*T)(\pi^*\tau) = \pi^*(T(\tau)) \quad .$$

It is easy to see that if T is surjective then π^*T is surjective as well. Also, π^* defines an isomorphism between $\text{Ker } T$ and $\text{Ker } (\pi^*T)$. Now suppose that Γ is *SEA*. Then we define h_{Γ_1} by $h_{\Gamma_1}(\phi) = h_{\Gamma}(\pi^*(\phi))$. By our observations above, h_{Γ_1} satisfies all the six axioms. ■

Proposition 2.2 *The unital free semigroup W_2 on two generators is not SEA.*

Proof: Suppose that our six axioms hold. The elements of W_2 can be identified with finite words on the two-letter alphabet $\{a, b\}$. The unit is the empty word E , the multiplication is the concatenation. Let $S = \{0, 1\}$, the cyclic group of two elements. Let $\phi_a \subset \Sigma(W_2, S)$ be defined the following way. Let $\tau \in \phi_a$ if and only if $\tau(\gamma) = 0$, when γ starts with the letter a . Then ϕ_a is a linear subshift. Similarly, we can define ϕ_b . There is an injective, continuous, W_2 -invariant map $T : \Sigma(W_2, S) \rightarrow \phi_a$, defined by $T(\tau)(\gamma) = \tau(b\gamma)$, $T(\tau)(E) = 0$ and $T(\tau)(a\gamma') = 0$ for any $\gamma' \in W_2$. By our axioms, $h(\phi_a) = 1$. Similarly, $h(\phi_b) = 1$. Notice that $\phi_a \cap \phi_b$ has two elements, hence $h(\phi_a \cap \phi_b) = 0$. Therefore,

$$1 = h(\Sigma(W_2, S)) = h(\phi_a) + h(\phi_b) = 2 \quad .$$

Thus we got a contradiction. ■

3 Groups of finite type

First we recall the notion of groups of finite type [2]. Let Γ be a finitely generated group, then we call it a group of finite type if there exists a finite simplicial complex $B\Gamma$, such that its universal cover $E\Gamma$ is contractible. It is important to remember that the homotopy type of $B\Gamma$ depends only on Γ , therefore it is meaningful to call the Euler-characteristic of Γ . We shall denote this integer by $e(\Gamma)$. Free groups, torsion free nilpotent groups and the

fundamental groups of negatively curved Riemann manifolds are all groups of finite type. Note that groups of finite type are of torsion-free and finitely presented. The free product of two groups of finite type is again a group of finite type. Now we can state the main result of our paper.

Theorem 1 *If Γ is a group of finite type and has nonzero Euler-characteristic, then Γ is not a SEA group.*

Before proving our Theorem it is worth to mention some related facts. By Stallings's Theorem, amenable groups of finite type always have zero Euler-characteristic. On the other hand free and surface groups have negative Euler-characteristic. Note that it is not enough to prove our Theorem for free groups and then apply Proposition 2.1. There exists groups with Kazhdan's Property (T), which are of finite type and nonzero Euler-characteristic. These groups have no free factors.

4 The Proof of the Theorem

Let $\widetilde{K}$ be an infinite, simplicial complex with a free and simplicial Γ -action such that the quotient space $K = \widetilde{K}/\Gamma$ is a finite simplicial complex. Note that $\widetilde{K}$ does not need to be a universal covering, just a simplicial covering of K with deck-transformation group Γ . We shall denote by $\widetilde{K}_i$ the set of i -dimensional simplices in $\widetilde{K}$ and by $C^i(\widetilde{K})$ is the group of $\mathbf{Z}_2$ -valued functions on $\widetilde{K}_i$, they are also known as the $\mathbf{Z}_2$ -valued i -cochains of $\widetilde{K}$.

Then we have the $\mathbf{Z}_2$ -cohomology complex as usual, where $\mathbf{Z}_2$ is the cyclic group of two elements.

$$C^0(\widetilde{K}) \xrightarrow{d_0} C^1(\widetilde{K}) \xrightarrow{d_1} C^2(\widetilde{K}) \xrightarrow{d_2} \dots \xrightarrow{d_{n-1}} C^n(\widetilde{K})$$

(here n is the largest dimension of a simplex in K). The definition of the coboundary maps d_i is very easy in this case. Simply, if $f \in C^i(\widetilde{K})$ and $\sigma \in \widetilde{K}_{i+1}$, then $df(\sigma) = \sum_{\tau_j} f(\tau_j)$, where the sum is taken over the i -dimensional faces of the simplex σ . Then we consider the $\mathbf{Z}_2$ -cohomologies as usual. $H^0(\widetilde{K}) = \text{Ker } d_0$, $H^1(\widetilde{K}) = \text{Ker } d_1 / \text{Im } d_0$, $H^2(\widetilde{K}) = \text{Ker } d_2 / \text{Im } d_1$ and so on, finally $H^n(\widetilde{K}) = C^n(\widetilde{K}) / \text{Im } d_{n-1}$. Now we use the entropy in order

to define some Betti numbers. First note that $C^i(\widetilde{K}) \cong \Sigma(\Gamma, S_i)$, where $S_i = \sum_{j=1}^{n_i} \mathbf{Z}_2$. Here n_i is the number of i -simplices in the quotient space K . Thus $h(C^i(\widetilde{K})) = \log_2 |S_i| = n_i$. Notice that all the kernels and images as well all the $C^i(\widetilde{K})$'s are linear subshifts. Thus the entropy betti numbers can be defined as follows : $b_h^0(\widetilde{K}) = h(\text{Ker } d_0)$, $b_h^1(\widetilde{K}) = h(\text{Ker } d_1) - h(\text{Im } d_0)$, $\dots$, $b_h^i(\widetilde{K}) = h(\text{Ker } d_i) - h(\text{Im } d_{i-1})$, $\dots$, $b_h^n(\widetilde{K}) = h(C^n(\widetilde{K})) - h(\text{Im } d_{n-1})$. One also have the entropy Euler-characteristic: $e_h(\widetilde{K}) = b_h^0(\widetilde{K}) - b_h^1(\widetilde{K}) + b_h^2(\widetilde{K}) - \dots + (-1)^n b_h^n(\widetilde{K})$.

Proposition 4.1 $e_h(\widetilde{K}) = e(K)$ (i.e. the usual topological Euler-characteristic of the finite simplicial quotient complex K).

Proof: We follow Cohen's proof for the analogous formula for the L^2 -Betti numbers [3]. He proved that $e(K) = e_{(2)}(\widetilde{K})$, where $e_{(2)}(\widetilde{K})$ denotes the L^2 -Euler characteristic. (this was proved in the differential setting by Atiyah himself in his classical Asterisque paper [1]). By our axioms we have the following $n + 1$ equations.

$$h(C^0(\widetilde{K})) = h(\text{Im } d_0) + h(\text{Ker } d_0)$$

$$h(C^1(\widetilde{K})) = h(\text{Ker } d_1) + h(\text{Im } d_1)$$

.

.

$$h(C^{n-1}(\widetilde{K})) = h(\text{Ker } d_{n-1}) + h(\text{Im } d_{n-1})$$

$$h(C^n(\widetilde{K})) = h(C^n(\widetilde{K}))$$

(The last row is not a misprint !) Now add up the equations above with alternating sign. This gives the proof of our proposition. ■

Now let us turn back to the proof of the Theorem. Let Γ be a group of finite type satisfying the *SEA* property. We need to prove that $e(\Gamma) = 0$. Consider the $B\Gamma$ finite simplicial classifying space. Since Γ is of finite type, the universal covering $\widetilde{K}$ is contractible. Hence, all the $\mathbf{Z}_2$ -cohomologies of $\widetilde{K}$ vanish, except the zeroth. Here, we have the constant functions representing the cohomology space. By our axioms, this linear subshift has zero

entropy. Consequently, all the Betti numbers are vanishing. Thus by our Proposition $e(\Gamma) = 0$. ■

References

- [1] M. F. Atiyah, *Elliptic operators, discrete groups and von Neumann algebras*, Asterisque, 32-33 (1976), pp. 43-72
- [2] K. Brown, *Cohomology of groups*, Graduate Texts in Mathematics , 87 (1982)
- [3] J. Cohen, *von Neumann dimension and the homology of covering spaces*, Quart. -J. -Math. -Oxford. -Ser(2) , 30 (1979) pp. 133-142
- [4] M. Coornaert, A. Papadopoulos, *Symbolic Dynamics and Hyperbolic Groups*, Lecture Notes in Mathematics, 1539 (1993)
- [5] S. Friedland, *Entropy of graphs, semigroups and groups*, in Ergodic Theory of $\mathbf{Z}^d$ -actions, Edited by M. Pollicott and K. Schmidt, (1996) pp. 319-343
- [6] J. Kieffer, *A generalized Shannon-McMillan theorem for the action of an amenable group on a probability space*, Annals of Probability 3 (1975), pp. 1031-1037
- [7] D. Lind, K. Schmidt and T. Ward, *Mahler measure and entropy for commuting automorphism of compact groups*, Inventiones Math. 101 (1990), pp. 593-629
- [8] T. Ward, Q. Zhang, *The Abramov-Rokhlin entropy addition formula for amenable group actions*, Monatsch. Math. 114 (1992) , pp. 317-329
- [9] S. A. Yuzvinskii, *Metric properties of endomorphisms of compact groups*, American Mathematical Society Translations, 66 (1968) pp. 63-98